Integrated Oyster Reef and Salt Marsh Restoration as an MEP Compliance Strategy Curtis S. Felix – Vice Chair, Wellfleet CWMPC cfelix@planktonpower.net



Water Quality Indicators

- Multiple data sources seem to agree that the inner harbor area exhibits:
 - Iow dissolved oxygen
 - Elevated chlorophyll a
 - Elevated nitrogen concentration
- Trends increase from south to north
- Mayo Creek outfall quality is low

Wellfleet Green Infrastructure Goals

- Protect and enhance the Wellfleet Harbor ecosystem now → think 1600's function
- Adaptive approach to nutrient management
- Identify low-cost, sustainable approaches
- Use best marine science focusing on
 - Integrated ecosystems
 - oyster reef restoration
 - salt marsh restoration

Program Elements

- 1. Oyster Propagation Project (2 acres) Wellfleet/UMass Boston/USDA/SPAT/MOP/Provincetown Center/Env. Partners
- 2. Oysterfest Shell Recycling NOAA/SPAT
- 3. Seaclam Cultch Program Wellfleet/USDA/MOP/SPAT
- 4. Townwide Shell Recycling DPW/Transfer Station Collection
- 5. Salt Marsh Restoration
 - Mayo Creek (20 acres)
 - Herring River (890 acres)
- 6. Composting toilets for marina nutrient removal
- 7. Traditional Needs Assessment
- 8. Townwide GIS Septic/Well Database

Feasibility

30 years of science documents the nitrogen removal capacities of oysters and salt marshes

Biodiversity of oyster reefs and salt marshes:
■ "habitat today → fish tomorrow"
Buffers shoreline erosion and ocean acidification

 Similar projects have been on-going in TX, NC, FL, Chesapeake Bay for over 15 years

Results in Maryland

MD: Governor O'Malley's Oyster Restoration and Aquaculture Development Plan 2009.

- increased Maryland's network of oyster sanctuaries from 9 percent to 24 percent;
- 2. increased leased areas for oyster aquaculture and streamlined the permitting process;
- 3. established a \$2.2 million financial assistance program for aquaculture interests; and
- 4. maintained 76 percent of the Bay's remaining habitat for targeted, sustainable, and scientifically managed public oyster fishery.
- Since implementation:
 - 28 new oyster farming leases have been approved on about 650 acres.
 - 52 lease applications covering 620 acres are currently being processed
 - MSX and Dermo have fallen to lowest levels ever recorded
 - Highest SPAT survival rates since 1985
 - Overall biomass up 44%
 - \$7.5 Million committed for 2013

http://www.oysterrecovery.org/Content/Content/1/Documents/2012 Octo ber3 PressRelease 2012 Oyster Planting Season Results.pdf



Outreach Effort

- Cape Cod Commission and DEP Staff
- Wellfleet Forum
- Shellfish Advisory Board/Planning Board/FinCom/ConsCom/Natural Resources Advisory
- Harbor Master/Health Department/Shellfish Department/DPW
- Board of Selectmen
- Cape Cod National Seashore
- Division of Marine Fisheries
- Non-Resident Taxpayer's Association
- WOMR/WHAT Theatre/Preservation Hall/LCAT/Newspapers
- Oysterfest

Results – Oyster Efforts

- 4 million new oysters in study area & 20% nitrogen reduction
- 50 million oysters from cultch programs' 1,200 tons of shell
- 3 billion gallons of added water filtration per day
- Pilot projects show start-up costs are minimal and maintenance is low (under \$15/lb of N removed)
- Anecdotal:
 - Huge influx of shrimp, tunicates, quahogs, crabs, snails, invertebrates, small fish
 - Significant reduction in mud level
 - Return of menhaden
 - Paired diamond back turtles foraging, heavy turtle use

Results – Oyster Efforts

Awards:


Mass Recycle – Municipal Innovation

November 2012

 American Council of Engineering Companies – Engineering Excellence Award

March 2013

Nitrogen Removal Smaller Scale Projects



Program Costs and Number of Human Nitrogen Equivalents Removed per Year


\$/Ib Nitrogen Removed^{1,2,3} People Equivalents³

1. Town Sea Clam Cultching '10 '11 '12	\$ 0.75	(\$50,000)	8,013
2. Mayo Creek Salt Marsh Restoration	\$ 1.78	(\$100,000)	336
3. Oysterfest Shell Recycling '11&'12	\$ 2.01	(\$2,000)	119
4. Oyster Propagation Zone	\$ 5.67	(\$50,000)	1,055
5. Herring River Salt Marsh Restoration	\$23.98	(\$60 million)	14,963
6. Baker Field Bathrooms	\$93.68	(\$324,000)	28
7. Sewering options (Cape Cod Commission)	\$500-\$1,000 (\$60 million)		450(CD Sewer)

- 1. Costs only; economic benefits dwarf costs in most cases;
- 2. Merrill/Cornwell 2002 Role of Oligohaline Marshes in Estuarine Nutrient Cycling
- 3. M Rice "Environmental Impacts of Shellfish Aquaculture: Filter feeding to Control Eutrophication"

Nitrogen cycling on oyster reefs

Denitrification, after Kellogg



Wellfleet Harbor Oyster Propagation Study Area



Site visit May 24, 2011 9 am



Fall 2011 after Cultch and Oysterfest

Notice thin white lines which are cultch visible in Google

Photo:

- Oysterfest oystershell (two small areas to the left represent 900,000 spat on shell
- Large cultched area represents 2 million oysters from natural set on seaclam shell and has increased to 4 million in 2012 © 2012 Google

Site visit July 16, 2011 9 am After initial cultch before receiving "set"


Site visit August 2011 after 1 month Example of spawn setting on cultch



Site visit September 2011 showing Even more set and good initial growth

Site visit October 17, 2012 8 am Showing significant 1 yr growth and another summer "set"





Why Recycle Shells?



Oyster, Clam, Mussel, Scallop only

Next generation of oysters will attach to this shell

Water Quality oysters consume algae filtering 50 gal/day

Critical Habitat for other fish "habitat today ... fish tomorrow"

Erosion control oysters form natural breakwaters that protect shoreline

Saving Money \$\$\$ no cost for out of Town disposal


Wellfleet Oysterfest 2012

- **25,000** visitors
- 100,000 oysters served
- 5.2 tons of shell recycled (NOAA Sponsored)
- 43% removed from 12.1 ton solid waste stream
- For every oyster eaten 6 were returned to the water or 577,448 oysters saved!
- 3 yr total= 15 tons of shell/ 900,000 SPAT






Oysterfest October 15, 2012 Can you find the 14 spat on this one oyster shell?



Success Rate



Number of Live Spat




Shellfish Constable Andy Koch with cultch spreader enroute to dropoff oysterfest shells

2 year olds attached to cultch otherwise not possible due to pure sand bottom



Wellfleet Oyster Propagation Project

Water Quality Monitoring Provincetown Center for Coastal Studies







Water Sampling Grid (not shown are 2 points further north in Duck Creek)



Total Nitrogen





< 21 [0.30] - Excellent 21-27 [0.3-0.39] - Good 27-35 [0.39-0.50] - Fair 35-50 [0.5-0.7] - Moderate Impairment 50-57 [0.7-0.8] - Significant Impairment > 57 [0.8] - Severe Degradation Water around oyster project is higher in nitrogen ranging from fair to severe Nitrogen Sink? @ degradation oyster densities 600-1,200 per m²

Total Nitrogen (uM [mg/L])